

COHERENT TRANSPORT THROUGH A QUANTUM TROMBONE.

N.K. Allsopp and C.J. Lambert,

School of Physics and Materials

Lancaster University

Lancaster LA1 4YB

U.K.

ABSTRACT.

We highlight a new “transverse” interference effect, arising from the coupling between electrons and holes, induced by a superconducting island in contact with a normal metal. As an example we compute the electrical conductance G of a T-shaped mesoscopic sample, formed by joining a horizontal bar of normal metal to a vertical leg of the same material. With a superconducting island located on the vertical leg and the current flowing horizontally, we examine the dependence of G on the distance L of the island from the horizontal bar. Of particular interest is the differential conductance $G(E)$ at an applied voltage $E = eV$, which due to quantum interference between electrons and Andreev reflected holes, is predicted to oscillate with both L and E . For a system with a spherical Fermi surface, the period of oscillation with E is $\pi E_F/k_F L$ and with L is $\pi E_F/k_F E$, where E_F is the Fermi energy and k_F the Fermi wavevector.

PACS Numbers. 72.10.Bg, 73.40.Gk, 74.50.

1. Introduction.

Mesoscopic superconductors, formed by adding one or more superconducting islands to a phase coherent normal host, represent a new class of quantum systems, whose realization has been made possible by recent advances in nano-lithographic techniques[1,2,3]. The possibility of such structures has led many theorists to consider incorporating superconductivity into modern theories of phase coherent transport[4-13] and several experimental groups to carry out related investigations[14-18]. In this paper, we analyze a simple demonstration experiment, carried out originally to address the prediction of superconductivity induced conductance suppression[12,13]. The experiment[18] measures the electrical conductance G of a T-shaped sample of size less than the quasi-particle phase breaking length, formed by joining a horizontal bar of silver to a vertical leg of the same material. With a superconducting island located on the vertical leg, a distance L from the T-junction and the current flowing horizontally, the dependence of G on the distance L and on the applied magnetic field was measured. As the applied field was lowered and the island allowed to become superconducting, macroscopic conductance changes of arbitrary sign were obtained, in agreement with the predictions of references[12,13].

In what follows, we compute the conductance $G(eV) = \partial I / \partial V$ of such a T-shaped “trombone” as the superconducting island is slid along the vertical leg. To our knowledge, this quantity has not yet been measured for such structures, although the work of references [14-18] demonstrates that $G(eV)$ is experimentally accessible. We note that at zero temperature, in the strict linear response limit of vanishing applied voltage, the conductance oscillates with L on the scale of the Fermi wavelength λ_F . This rapid variation arises from normal reflection at the interface with the island and is found in quantum modulated transistors[19,20,21], with no superconductivity. In contrast, we predict that the differential conductance at finite voltage V is sensitive to quantum interference between electrons and Andreev reflected holes and show that this oscillates

with L on the much longer scale $\sim \pi E_F/(Ek_F)$, where $E = eV$, E_F is the Fermi energy and k_F the Fermi wavevector. Such an effect is reminiscent of Tomasch oscillations in tunnel junctions[22,23], although in the present context, the current flow is transverse to the direction of the geometric feature causing interference.

2. Calculation of the electrical conductance.

To demonstrate this new interference effect, one notes that if the ends of the horizontal bar of the trombone are connected to external reservoirs by normal, current carrying leads, then the two probe, differential electrical conductance is given by

$$G(E) = e[\partial I/\partial(\mu_1 - \mu_2)]_{(\mu_1 - \mu_2 = E)},$$

where μ_1, μ_2 are the chemical potentials of the reservoirs. At zero temperature, the analysis of references [7,8], yields, in units of $2e^2/h$,

$$G(E) = \frac{1 - R_{pp}(E) + R_{hp}(E) + \alpha(T_{hh'}(E) - T_{ph'}(E))}{1 + \alpha}, \quad (1)$$

where

$$\alpha = \frac{R_{hp}(E) + T_{h'p}(E)}{R_{p'h'}(E) + T_{ph'}(E)}$$

and the notation of reference[7] has been adopted. The coefficients $R_{pp}(E), R_{hp}(E)$ ($R_{h'h'}(E), R_{p'h'}(E)$) are probabilities for normal and Andreev reflection respectively for quasi-particles (quasi-holes) from the left (right) reservoir, while $T_{p'p}(E), T_{h'p}(E)$ ($T_{hh'}(E), T_{ph'}(E)$) are the corresponding transmission probabilities. These are obtained by tracing over combinations of sub-matrices of the scattering matrix of the structure[7,8].

Consider first the T-shaped structure shown in figure 1, which comprises three one-dimensional wires, connected at a node. In the absence of superconductivity, the Bogoliubov - de Gennes equation

$$\begin{pmatrix} H_0(\underline{r}) & \Delta(\underline{r}) \\ \Delta^*(\underline{r}) & -H_0^*(\underline{r}) \end{pmatrix} \begin{pmatrix} \psi(\underline{r}) \\ \phi(\underline{r}) \end{pmatrix} = E \begin{pmatrix} \psi(\underline{r}) \\ \phi(\underline{r}) \end{pmatrix} \quad (2),$$

reduces to two separate equations for particles and holes. For clean wires, in a region occupied by the j th wire, a solution of the particle (hole) equation consists of incoming and outgoing plane waves of amplitudes $v_k^{-1/2}A_j, v_k^{-1/2}B_j$ ($v_q^{-1/2}\bar{A}_j, v_q^{-1/2}\bar{B}_j$) respectively, where v_k (v_q) is the group velocity of a particle (hole) of energy $E = \hbar^2 k^2/2m - E_F = E_F - \hbar^2 q^2/2m$. These satisfy equations of the form

$$\begin{pmatrix} B_1 \\ B_2 \\ B_3 \end{pmatrix} = S(E) \begin{pmatrix} A_1 \\ A_2 \\ A_3 \end{pmatrix} \quad (3)$$

and

$$\begin{pmatrix} \bar{B}_1 \\ \bar{B}_2 \\ \bar{B}_3 \end{pmatrix} = S^*(-E) \begin{pmatrix} \bar{A}_1 \\ \bar{A}_2 \\ \bar{A}_3 \end{pmatrix} \quad (4).$$

To investigate the possibility of geometric resonances, we now introduce a superconducting boundary, a distance L along the vertical leg of the structure. For convenience, we consider an ideal boundary, with negligible normal scattering, which induces coupling between particles and holes only. If $r_a(E)$ ($\bar{r}_a(E)$) are Andreev reflection coefficients for particles (holes) incident on the interface, then the coupling takes the form

$$\bar{A}_3 = r_a(E)B_3 \quad A_3 = \bar{r}_a(E)\bar{B}_3 \quad (5).$$

This pair of equations allows us to eliminate the amplitudes $A_3, \bar{A}_3$ from equations (3) and (4), to yield an equation of the form (see appendix)

$$\begin{pmatrix} B_1 \\ B_2 \\ \bar{B}_1 \\ \bar{B}_2 \end{pmatrix} = \begin{pmatrix} S^{pp}(E) & S^{ph}(E) \\ S^{hp}(E) & S^{hh}(E) \end{pmatrix} \begin{pmatrix} A_1 \\ A_2 \\ \bar{A}_1 \\ \bar{A}_2 \end{pmatrix} \quad (6),$$

where the 2x2 sub-matrices $S^{ij}(E)$ satisfy the following particle-hole symmetry relations $S^{hh}(-E) = [S^{pp}(E)]^*$ and $S^{ph}(E) = -[S^{hp}(-E)]^*$. All reflection and transmission coefficients are obtained from the modulus squared of elements of these sub-matrices. In particular, $T_{p'p}(E) = |S_{21}^{pp}|^2$, $R_{hp}(E) = |S_{11}^{hp}|^2$, $T_{hh'}(E) = |S_{12}^{hh}|^2$ and $R_{p'h'}(E) = |S_{22}^{ph}|^2$.

As an example, consider a spatially symmetric scatterer, for which

$$S(E) = \begin{pmatrix} r & t & t \\ t & r & t \\ t & t & r \end{pmatrix}. \quad (7)$$

In this case, the conduction formula (1) reduces to $G(E) = (R_{ph}(E) + R_{h'p'}(E) + T_{hh'}(E) + T_{p'p}(E))/2$. and in the limit $E/E_F \ll 1$, one obtains (see appendix)

$$T_{p'p}(E) \simeq |t|^2 |1 + r^* t/d|^2 \quad (8),$$

$$T_{hh'}(E) \simeq |t|^2 |1 + r t^*/d|^2 \quad (9),$$

$$R_{hp} \simeq R_{p'h'} \simeq |t|^4 / |d|^2 \quad (10),$$

where

$$d \simeq -|r|^2 - \exp[-2i(k - q)L] \quad (11).$$

In the above expressions, only the denominator d depends on the length L . At zero energy, since $k - q \simeq k_F E/E_F$, this dependence vanishes. At finite energy, there is a slow oscillation with period given by

$$k_F L E/E_F = \pi \quad (12).$$

3. Numerical results in two-dimensions.

The above analysis demonstrates that transverse geometric oscillations are a generic feature of structures of the kind shown in figure 1. For a more realistic interface, which induces both normal and Andreev reflection, it is clear from the analysis of references[19-21] that there would be an additional, more rapid oscillation on the scale of $k_F L$. To illustrate this feature and to show that slow oscillations survive in more realistic systems of finite width, we now present the results of a numerical simulation of the 2-dimensional T-shaped structure, shown in figure 2 and described by a tight binding Bogoliubov - de Gennes operator of the form

$$H = \begin{pmatrix} H_0 & \Delta \\ \Delta^* & -H_0^* \end{pmatrix} \quad (13).$$

In this equation H_0 a nearest neighbour Anderson model on a square lattice, with off-diagonal hopping elements of magnitude unity and Δ a diagonal matrix with on site, particle-hole couplings. The latter are chosen to vanish in the normal regions and to have a value Δ_0 in the region occupied by the superconducting island (shown black in figure 2). The scattering region is chosen to be $M = 30$ sites wide and 6 sites long and is connected to external leads of width M . Within the leads, the diagonal elements are equal to ϵ_0 and within the insulating portion of the scattering region (shown shaded) the diagonal elements are set to an arbitrarily large value $\bar{\epsilon}$. Within the remaining (conducting) portion of the scattering region, diagonal elements of H_0 are chosen to be random numbers, uniformly distributed between $\epsilon_0 - W$ and $\epsilon_0 + W$. In what follows, the choice $\epsilon_0 = 0$, $\bar{\epsilon} = 50$, $\gamma = 1$ was made and free end boundary conditions employed. In the experiments of reference[18], the conductance per channel was estimated to be of order $0.1 e^2/h$, which for the system of figure 2, is obtained with the choice $W \simeq 0.1$. The results of figure 3 were obtained with this choice of W and $\Delta_0 = 0.2$.

As well as possible oscillatory dependencies on E and L , it is of interest to examine the effect of a magnetic field. Therefore for each set of random diagonal elements, the conductance $G(E)$ was first obtained with zero magnetic field, then recomputed with a finite flux ϕ through the conducting region. For each choice of E and L , the average conductance $G(E, L)$ was computed from results obtained for 100 different realisations of the random diagonal elements, with the same sets of random numbers being used for each E and L . The squares in figure 3 show results for $\phi = 0$, while the circles show results for $\phi = 6\phi_0$, where $\phi_0 = hc/e$. For the latter, the field was introduced through a Peierls substitution in the off-diagonal elements of H_0 .

At zero energy, the conductance oscillates on the scale of the lattice spacing, which for the system of figure 2 is of order the Fermi wavelength λ_F . At this energy, particle-hole symmetry ensures that there is no interference between electrons and Andreev reflected holes from the normal-superconducting interface. At finite energies, the wavelengths

of particle and holes differ by an amount proportional to E and therefore quantum interference on a length scale proportional to E^{-1} can occur. This new oscillatory effect, is distinct from more familiar Tomasch[22] and Macmillan-Rowell[23] oscillations, since the latter occur in tunnel junctions, while in the structure considered here, there is no insulating barrier. In addition, the effect reported here occurs for $E < \Delta_0$ as well as $E > \Delta_0$.

To obtain a crude estimate of the periods of oscillation shown in figure 3, one notes that in the absence of disorder and superconductivity, a vertical, normal lattice of width two sites possesses a dispersion relation for particles of the form $E = -2\gamma[\cos k_y a + \cos k_x^n a]$ and for holes, $E = +2\gamma[\cos q_y a + \cos k_x^n a]$, where $k_x^n a = n\pi/3$, $n = 1, 2$. Choosing $E = 0$, yields values for the allowed wavevectors k_F^n on the Fermi surface given by $k_F^n a = \pi/3$ or $3\pi/3$. Hence for small E , $E \simeq \gamma \sin k_F^n a (k_y - q_y) a = \gamma(k_x - q_x) a \sqrt{3}/2$. and the period of oscillation obtained from equation (12) is given by

$$(L/a)(E/\gamma) \simeq 2\pi/\sqrt{3} \quad (14).$$

While this is a somewhat crude estimate, one notes that for $E/\gamma = 0.2$, this yields $L \simeq 18$, which is of the same order as the period shown in the bottom graph of figure 3.

Figure 3 shows that oscillations occur both in zero and finite magnetic fields and as a consequence, the magneto-conductance can change sign, if either L or E are varied. The oscillations predicted by figure 3 are a finite fraction of the total conductance and therefore should be observable experimentally. In practice it is not possible to slide the superconducting island of a quantum trombone, without growing a separate sample with an independent set of random impurities. Consequently oscillations on the scale of λ_F are difficult to observe and would be measured as random sample to sample fluctuations. In contrast the longer scale variation in the differential conductance should be accessible experimentally. Indeed, by fixing the length L and varying E , the oscillations reported above should be observable in a single sample. For $L = 11$, this behaviour is illustrated

in figure 4, which shows a slow variation with E on a scale proportional to L^{-1} . For $L = 11$, equation (14) yields a the period of $E/\gamma \simeq 0.3$, compared with a value $E/\gamma \simeq 0.2$ obtained from figure 4.

4. Discussion.

We have demonstrated that a superconducting island in contact with a normal mesoscopic conductor, produces geometric resonances, even if the superconductor is far removed from the current path. Such effects should be observable in the mesoscopic structures described in reference [18]. The above results, which are based on the assumption of a rigid order parameter, do not explain a weak magnetic field feature found in [18]. Figures 3 and 4 also show that the magneto-conductance can change sign with varying E and L , but the magnitude of the effect is much smaller than that observed in reference [18]. Since the theory of reference[12] predicts that the electrical conductance of a hybrid superconducting structure is sensitive to small changes in the magnitude of the superconducting order parameter, a possible explanation of the measured small field feature is that weak proximity tails present in the silver are modified by the applied field. To avoid such tails, it would be of interest to repeat these experiments, but with a magnetic normal host such as Nickel[24]

Appendix.

Combining equations (3) and (5) yields $B_3 = S_{33}A_3 + S_{31}A_1 + S_{32}A_2 = \bar{A}_3/r_a(E)$. Similarly combining equations (4) and (5) yields $\bar{B}_3 = \bar{S}_{33}\bar{A}_3 + \bar{S}_{31}\bar{A}_1 + \bar{S}_{32}\bar{A}_2 = A_3/\bar{r}_a(E)$, where the i, j th element of $S_{ij}(E)$ has been denoted S_{ij} and the i, j th element of $S_{ij}^*(-E)$ denoted $\bar{S}_{ij}$. These yield expressions for $A_3, \bar{A}_3$ of the form

$$A_3 = \alpha_1 A_1 + \alpha_2 A_2 + \beta_1 \bar{A}_1 + \beta_2 \bar{A}_2$$

and

$$\bar{A}_3 = \bar{\beta}_1 A_1 + \bar{\beta}_2 A_2 + \bar{\alpha}_1 \bar{A}_1 + \bar{\alpha}_2 \bar{A}_2,$$

where

$$\alpha_1 = \bar{S}_{33}S_{31}/d, \alpha_2 = \bar{S}_{33}S_{32}/d,$$

$$\bar{\alpha}_1 = S_{33}\bar{S}_{31}/d, \bar{\alpha}_2 = S_{33}\bar{S}_{32}/d,$$

$$\beta_1 = \bar{S}_{31}/(d r_a(E)), \beta_2 = \bar{S}_{32}/(d r_a(E)),$$

$$\bar{\beta}_1 = S_{31}/(d \bar{r}_a(E)), \bar{\beta}_2 = S_{32}/(d \bar{r}_a(E)), \text{ with } d = (r_a(E)\bar{r}_a(E))^{-1} - S_{33}\bar{S}_{33}.$$

Substituting these back into equations (3) and (4) yields equation (6), with

$$S^{pp} = \begin{pmatrix} S_{11} + S_{13}\alpha_1 & S_{12} + S_{13}\alpha_2 \\ S_{21} + S_{23}\alpha_1 & S_{22} + S_{23}\alpha_2 \end{pmatrix}$$

$$S^{ph} = \begin{pmatrix} S_{13}\beta_1 & S_{13}\beta_2 \\ S_{23}\beta_1 & S_{23}\beta_2 \end{pmatrix}$$

$$S^{hh} = \begin{pmatrix} \bar{S}_{11} + \bar{S}_{13}\bar{\alpha}_1 & \bar{S}_{12} + \bar{S}_{13}\bar{\alpha}_2 \\ \bar{S}_{21} + \bar{S}_{23}\bar{\alpha}_1 & \bar{S}_{22} + \bar{S}_{23}\bar{\alpha}_2 \end{pmatrix}$$

$$S^{hp} = \begin{pmatrix} \bar{S}_{13}\bar{\beta}_1 & \bar{S}_{13}\bar{\beta}_2 \\ \bar{S}_{23}\bar{\beta}_1 & \bar{S}_{23}\bar{\beta}_2 \end{pmatrix}$$

For a N-S interface at a point x_0 along the x -axis, Andreev's approximation for reflection amplitudes is $r_a(E) = \exp i[-\theta - \phi + (k - q)x_0]$, $\bar{r}_a(E) = \exp i[-\theta + \phi + (k - q)x_0]$,

where ϕ is the phase of the superconducting order parameter and $\theta = \tan^{-1}[(|\Delta|^2 - E^2)^{1/2}/E] + \pi\Theta(-E)$. In view of particle-hole symmetry $r_a(E) = -\bar{r}_a^*(-E)$, as expected.

For a symmetric scatterer, with S-matrices of the form

$$S(E) = \begin{pmatrix} r & t & t \\ t & r & t \\ t & t & r \end{pmatrix}, \quad \bar{S}(E) = S^*(-E) = \begin{pmatrix} \bar{r} & \bar{t} & \bar{t} \\ \bar{t} & \bar{r} & \bar{t} \\ \bar{t} & \bar{t} & \bar{r} \end{pmatrix},$$

one obtains

$$T_{p'p}(E) = |t|^2 |1 + \alpha_1|^2,$$

$$T_{hh'}(E) = |\bar{t}|^2 |1 + \bar{\alpha}_2|^2,$$

$$R_{hp}(E) = |\bar{t}|^2 |\bar{\beta}_1|^2$$

and

$$R_{p'h'}(E) = |t|^2 |\beta_2|^2.$$

In these expressions,

$$\alpha_1 = \bar{t}/d\bar{\alpha}_2 = r\bar{t}/d$$

$$\bar{\beta}_1 = t/(d\bar{r}_a)\beta_2 = \bar{t}/(dr_a)$$

and

$$d = (r_a(E)\bar{r}_a(E))^{-1} - |r|^2.$$

For small E , provided t, r are slowly varying compared with $(E/E_F)k_F L$, one may evaluate r, t at $E = 0$ and write $\bar{t} \simeq t^*$ and $\bar{r} \simeq r^*$. This yields equations (8) to (11) of the main text.

Acknowledgements.

This work is supported by the SERC, the EC Human Capital and Mobility Programme and the MOD. It has benefited from useful conversations with V. Petrashov, V.C. Hui and F. Sols.

Figure Captions.

Figure 1. A quantum trombone formed from three one-dimensional wires connected at a node. This figure shows the arrangement of incoming and outgoing amplitudes for particles.

Figure 2. A quantum trombone formed from a two dimensional, tight-binding lattice of sites. The superconducting island is coloured black and located on the vertical arm of the trombone. The shaded area represents an insulating portion of the scattering region, while white area shows the weakly disordered conducting region. The current flows from left to right along the horizontal arm of the T-junction. The external leads and scatterer are of width $M = 30$ sites. The horizontal arm is 6 sites long. The horizontal arm and vertical leg are each 2 sites wide.

Figure 3. Results for the ensemble averaged differential conductance $G(E, L)$ as a function of the position L of the superconducting island, for $E = 0$ (top graph), $E = 0.1\gamma$ (middle graph) and $E = 0.2\gamma$ (bottom graph). In each case $\Delta_0 = 0.2$. The squares show results for zero applied magnetic field and the circles for an applied flux of $6\phi_0$ passing through the normal part of the sample, (ie. the white plus shaded regions of figure 2), where $\phi_0 = hc/e$.

Figure 4. As for figure 3, except that $G(E, L)$ is plotted as a function of E . In this figure $\Delta_0 = 0.2$ and $L = 11$.

References.

1. V.T. Petrashov and V.N. Antonov, Sov. Phys. JETP Lett. **54** 241, (1991)
2. V.T. Petrashov, V.N. Antonov and Persson, Physica Scripta, **T42** 136 (1992),
3. V.T. Petrashov, V.N. Antonov, P. Delsing and T. Claeson, Phys. Rev. Lett. **70** 347 (1993).
4. G.E. Blonder, M. Tinkham and T.M. Klapwijk, Phys. Rev. B. **25** 4515 (1982).
5. H. van Houten and C.W.J. Beenakker, Physica B. **175** 187 (1991).
6. B.Z. Spivak and D.E. Khmel'nitskii, Sov. Phys. JETP Lett, **35** 413 (1982)
7. C.J. Lambert, J. Phys.: Condens. Matter, **3** 6579 (1991)
8. C.J. Lambert, V.C. Hui and S.J. Robinson, J.Phys.: Condens. Matter, **5** 4187 (1993)
9. Y. Takane and H. Ebisawa, J. Phys. Soc. Jap. **61** 1685 (1992).
10. C.W.J. Beenakker, R.A. Jalabert and I.K. Marmorkos, Phys. Rev. **B48** 2811 (1993)
11. V.C. Hui and C.J. Lambert, Euro. Phys. Lett., **23** 302 (1993)
12. C.J. Lambert and V.C. Hui, J. Phys.: Condens. Matter **5** L651 (1993)
13. C.J. Lambert and V.C. Hui, Proc. 20th Int. Conf. on Low Temp. Phys. (1994) to be published
14. Y.W. Kwong, K. Lin, P.J. Hakonen, M.S. Isaacson and J.M. Parpia, Phys. Rev. B **44** 462 (1991).
15. P. Santhanam, C.C. Chi, S.J. Wind, M.J. Brady and J.J. Bucchignano, Phys. Rev. Lett. **66** 2254 (1991).
16. A. Kastalasky, A.W. Kleinsasser, L.H. Greene, R. Bhat and J.P. Harbison, Phys. Rev. Lett. **67** 3026 (1991)
17. D.R. Heslinga and T.M. Klapwijk, Phys. Rev. **B47** 5157 (1993)
18. V.T. Petrashov, V.N. Antonov, S.V. Maksimov, Sov. Phys. JETP Lett., **58** 49 (1993)
19. F. Sols, M. Macucci, U. Ravaioli and K. Hess, J. Appl. Phys. **66** 3892 (1988)

- 20. F. Sols, M. Macucci, U. Ravaioli and K. Hess, Appl. Phys. Lett. **54** 350 (1989)
- 21. H.U. Baranger, A.D. Stone and D.P. DiVincenzo, Phys. Rev. **B37** 6521 (1988)
- 22. W.J. Tomasch, Phys Rev Lett., **15** 672 (1965)
- 23. W.L. Macmillan and J.M. Rowell, Phys Rev Lett., **15** 453 (1966)
- 24. V.T. Petrashov Private communication.